

**Statewide Quality Advisory Committee
2014 Nominated Quality Measures**

SQAC Staff Recommends Evaluation of the Following Nominated Measures, by Focus Area

End-of-Life Care

1. CARE - Consumer Assessments and Reports of End of Life (NQF 1632)
2. Advance Care Plan (NQF 326)
3. Proportion admitted to hospice for less than 3 days (NQF 216)
4. Beliefs/Values Addressed (if desired by the patient) (NQF 1647)
5. Family Evaluation of Palliative Care
6. Family Evaluation of Hospice Care (NQF 208)
7. Palliative and End of Life Care: Dyspnea Screening and Management
8. Hospice Set: Pain Screening (NQF 1634)
9. Hospice Set: Pain Assessment (NQF 1637)
10. Hospice Set: Dyspnea Screening (NQF 1639)
11. Hospice Set: Dyspnea Treatment (NQF 1638)
12. Hospice Set: Patients Treated with an Opioid who are Given a Bowel Regimen (NQF 1617)

Patient-Centered Care

13. Use and Quality of Shared Decision-Making
14. Active Patient Engagement

Pediatric Care

15. Maternal Depression Screening (NQF 1401)
16. Developmental Screening in first 3 years of life (NQF 1448)
17. Depression screening by 18 years of age (NQF 1515)
18. Diagnosis of ADHD in primary care for school-aged children and adolescents (NQF 106)
19. PC-02 Cesarean Section (NQF 471)

Behavioral Health

20. Post discharge continuing care plan transmitted to next level of care provider upon discharge (HBIPS 7)
21. Post discharge continuing care plan created (HBIPS 6)
22. Patient Health Questionnaire: the PHQ-9 (NQF 712, 711, 710)
23. Alcohol Use Disorder Identification Test (AUDIT) (NQF 2152)
24. Generalized Anxiety Disorder 7-item (GAD-7)
25. Columbia Suicide Severity Rating Scale (C-SSRS)

Measures for Discussion – Do Not Meet a Focus Area

Staff Recommends Evaluation

Patient Safety

26. Patient Safety Composite (PSI 90) (NQF 531)
27. Catheter-Associated Urinary Tract Infections (NQF 138)
28. SSI Surgical Site Infection: SSI colon, SSI-abdominal hysterectomy (NQF 753)
29. Central-Line Associated Bloodstream Infection (NQF 139)
30. Hospital-onset methicillin resistant staphylococcus bacteremia aureus (MRSA) (NQF 1716)
31. Hospital-onset *C. difficile* (NQF 1717)

Leapfrog

32. Leapfrog measure set

CMS Hospital Measures

33. Pneumonia 30-day mortality rate (NQF 468)
34. Heart failure 30-day mortality rate (NQF 229)
35. AMI 30-day mortality rate (NQF 230)

Staff Recommends Further Discussion

CMS Hospital Measures

36. Fibrinolytic therapy received within 30 minutes of emergency department arrival (Hosp OP-2)
37. Aspirin at arrival (Hosp OP-4)
38. Timing of antibiotic prophylaxis (Hosp OP-6)
39. Prophylactic antibiotic selection for surgical patients (Hosp OP-7)
40. Immunization for influenza (IMM-2)

Primary Care Utilization

41. Total number of primary care visits
42. Total number of patients (one or more visits during report period)
43. Total number of primary care visits by asthma patients (5-64) as defined by HEDIS
44. Total number of primary care visits by diabetic patients, as defined by HEDIS
45. Total number of primary care visits by hypertension patients, as defined by HEDIS